

Why Create Healthy Communities? Research Evidence

Brian E. Saelens, Ph.D. Seattle Children's Hospital University of Washington

Snohomish County Healthy Communities by Design for All Ages
September 2009

Socio-ecological model of obesity





Evidence on Children & Healthy Communities

NIK Study Design

		Physical Activity Environment			
		High	Low		
Nutrition	High	High PAE/High NE environment - high walkable - good park availability/quality - healthy food environment	Low PAE/High NE environment - low walkable - poor park availability/quality - healthy food environment		
environment	Low	High PAE/Low NE environment - high walkable - good park availability/quality - unhealthy food environment	Low PAE/Low NE environment - low walkable - poor park availability/quality - unhealthy food environment		

Combining Walkability & Park Quality

- High physical activity environment
 High PAE = high walkable + high quality parks
- Low physical activity environment Low PAE = low walkable + no parks













Neighborhood Nutrition Evaluation


- Food establishments within and within ½ mile buffer of block group
- High nutrition
 - At least one supermarket and
 - 15 or fewer fast food establishments
- Low nutrition
 - No supermarket

OR

- A supermarket but
- 16 or more fast food establishments




Child BMI z-score (By Neighborhood Type)



- **Low PAE, Low NE**
- **Low PAE, High NE**
- **■** High PAE, Low NE
- **High PAE, High NE**


Child Overweight and Obesity (By Neighborhood Type)




- **Low PAE, Low NE**
- **Low PAE, High NE**
- High PAE, Low NE
- High PAE, High NE

Evidence on Adults & Healthy Communities

NQLS Neighborhood Categories




NQLS Study: Self-Reported Physical Activity



^{*} Adjusted for neighborhood clustering, gender, age, education, ethnicity, # motor vehicles/adult in household, site, marital status, number of people in household, and length of time at current address.


NQLS Study: Self-Reported Physical Activity



Adjusted for 'reasons for moving', sex, age, education, ethnicity, marital status, vehicles in household; neighborhood clustering

Sallis et al 2009 SSM

Body Mass Index (BMI)



Adjusted for age and sex

Built Environment - Walking Associations among Published 2005/2006 articles (Adults)

	Transport walking	Recreation walking
Density	+ (80%)	? (25%)
Distance to destinations	- (78%)	? (20%)
Land-use mix	+ (85%)	? (43%)
Street connectivity	? (50%)	? (0%)
Pedestrian infrastructure	? (25%)	+ (67%)
Traffic	? (25%)	? (14%)
Personal safety	? (43%)	? (43%)
Parks/open space	? (40%)	? (0%)
Aesthetics	? (17%)	+ (100%)

Evidence-based Recommendations

- Implement strategies that bring people closer to desired destinations
- Change the balance of facilitators/barriers to active transportation
 - Improve walking/biking infrastructure
 - Disincentivize driving alone
- Increase healthy food access while addressing costs
- http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5 807a1.htm for Community Strategies and Measurements to Prevent Obesity